

Adult Health in Great Britain, 2013



Coverage: **GB**

Date: **19 March 2015**

Geographical Area: **Region**

Theme: **Health and Social Care**


Key points

- In 2013 more than one in three adults (36%) reported having a long-standing illness or disability (LSI) and one in five (20%) reported having a limiting LSI.
- The likelihood that someone reported having an LSI was closely associated with age. In 2013 69% of people aged 75 and over reported having an LSI. This compared with 15% of people aged 16 to 24.
- People living in Wales (27%) were more likely to report having a limiting LSI than those living in either England (19%) or Scotland (20%).
- Cigarette smokers (37%) were more likely to report an LSI than those who had never smoked cigarettes (32%).

Summary

This report looks at self-reported long-standing illnesses and disabilities (LSIs) and self-reported LSIs that limit activities. A long-standing illness or disability (LSI) is anything that someone has considered to have troubled them over a period of time, or that they believe is likely to affect them over a period of time.

In 2013, more than one in three adults in Great Britain (36%) reported having a long-standing illness or disability. This had increased slightly compared with 2012 (34%) but was in line with the levels seen over the 2005 to 2012 period, Figure 1. One in five adults (20%) said that they had a limiting LSI. Although this had fallen slightly since 2005 (22%), there has been little change compared with 2012 (19%).

Figure 1: Self-reported long-standing illness or disability, Great Britain, 2005-2013

Source: Opinions and Lifestyle Survey, General Lifestyle Survey, General Household Survey - Office for National Statistics

Download chart

[XLS](#) [XLS format](#)
(27.5 Kb)

As expected, older adults were more likely to have reported having an LSI or limiting LSI than those in younger age groups. The likelihood of reporting an LSI is complex and related to a number of other factors. Unemployed people, people on low incomes, cigarette smokers and people living in Wales were more likely to have reported a limiting LSI than others. Within the group of cigarette smokers, higher consumption levels were associated with higher rates of self-reported LSI and limiting LSI.

The association between relationship status and self-reported LSI was complex. People who were married, widowed, divorced or separated reported higher rates of LSI than those who were single or cohabiting. However, age was the driving factor behind these differences. Once age and other factors were taken into account, single people were most likely to have reported having an LSI.

The most commonly reported difficulties caused by long-standing illness were with mobility; stamina breathing and fatigue; and dexterity. There were some differences between men and women in the reported effects of LSI.

Why do these results matter?


More than one in three people in Great Britain live with a long-standing illness or disability. This can affect many areas of a person's life, including [employment, relationships, education and the ability to live independently](#).

With the [number of older people in the UK increasing](#), long-standing illness and disability is an increasingly important issue for governments and health and social care providers. Long-term conditions account for [70% of health and social care spending and 50% of all GP consultations](#) in England with the [NHS Mandate](#) setting out what NHS England must do to improve the care of people with long-term conditions. The [Welsh Government](#) has a 10 year vision highlighting the need to improve management of long-term conditions in order to provide sustainable and effective health and social care services. NHS Scotland have set out a [National Action Plan](#) highlighting inequalities in health relating to deprivation and lifestyle factors and outlining an integrated approach to improve the quality of care for people with long-term conditions.

Self-reported long-standing illness or disability

Self-reported long-standing illness or disability (LSI), age and sex

The likelihood of reporting an LSI increased with age, Figure 2. Those aged 75 and over were more than four times as likely to have reported an LSI than those aged 16 to 24 (69% vs. 15%). They were also nearly six times as likely to have reported a limiting LSI (47% vs. 8%).

Figure 2: Self-reported long-standing illness or disability, by age, Great Britain, 2013


Source: Opinions and Lifestyle Survey - Office for National Statistics

Download chart

[XLS](#) [XLS format](#)

(27 Kb)

Overall, men reported lower levels of LSIs (34%) and limiting LSIs (19%) than women (37% and 21%). This continued the trend seen in recent years, Figure 3. However, health is linked with a number of other factors including age, employment status and economic activity. Once the effects of other factors were accounted for, there was no difference in the rates of LSI and limiting LSI between men and women.

Figure 3: Self-reported long-standing illness or disability, by sex, Great Britain, 2005-2013

Source: Opinions and Lifestyle Survey, General Lifestyle Survey, General Household Survey - Office for National Statistics


Download chart

[XLS](#) [XLS format](#)
(28 Kb)

Work and self-reported long-standing illness or disability

Unemployed people are defined as those who are not working, but are looking for work. Around one in three people who were unemployed (33%) reported having an LSI, compared with around one in four people in employment (24%). Unemployed people were more than twice as likely as people in employment to have reported a limiting LSI (20% vs. 9%), Figure 4. This reflects findings from the [Labour Force Survey](#), which show that during 2013, the unemployment rate was higher among those who had a disability.

Figure 4: Self-reported long-standing illness or disability, by employment status, Great Britain, 2013



Source: Opinions and Lifestyle Survey - Office for National Statistics

Notes:

1. International Labour Organisation (ILO) definitions of employment status have been used.
2. Unemployed people are those who are not currently in work, but are looking for work.

Download chart


[XLS](#) [XLS format](#)
(28 Kb)

Income and self-reported long-standing illness or disability

Low incomes were associated with higher rates of self-reported LSI. People with an income under £10,000 were nearly twice as likely to have reported an LSI (45%) than those with an income over £50,000 (24%), Figure 5.

Low income was also related to higher reports of limiting LSI and this remained the case once the effect of factors such as region, smoking and employment status had been removed. People with an income under £10,000 were more than three times as likely to have reported a limiting LSI (30%) than those who earned over £30,000 (9%). However, for those belonging to income groups beyond £30,000 a year there was little difference in the rates of self-reported limiting LSIs.

Figure 5: Self-reported long-standing illness or disability and gross annual personal income, Great Britain, 2013



Source: Opinions and Lifestyle Survey - Office for National Statistics

Notes:

1. Gross annual personal income covers all personal income before deductions for tax, National Insurance etc. It relates to income that is directly received (such as pay, benefits or interest from savings) and does not include income from a third party (such as a spouse or partner).

Download chart

[XLS](#) [XLS format](#)

(28 Kb)

Relationships and self-reported long-standing illness or disability


Reports of LSI were higher for those who were married (38%) or widowed, divorced or separated (54%) than people who were single (26%) or cohabiting (25%). However, age is the driving factor behind these differences. Those who are divorced, widowed or separated and, to a lesser extent those who are married, are older on average than those who are cohabiting or single. Once the effects of factors such as age and employment status were removed, single people were between 20% and 30% more likely to report a long-standing illness or disability than those in other relationship groups.

A similar pattern was seen in the association between relationship status and limiting LSI. More than twice as many people who were widowed, divorced or separated reported having a limiting LSI (35%) than single people (15%). However, once other factors such as age were accounted for, single people were 28% more likely to have reported a limiting long-standing illness or disability than people who were married or cohabiting and there was no clear difference between single people and people who were widowed, divorced or separated.

Cigarette smoking and self-reported long-standing illness or disability

Nearly a third of adults (32%) who had never smoked cigarettes reported having an LSI. This was lower than the proportion of cigarette smokers (37%) and ex-smokers (45%), Figure 6.

Figure 6: Cigarette smoking and self-reported long-standing illness or disability, Great Britain, 2013



Source: Opinions and Lifestyle Survey - Office for National Statistics

Notes:

1. Light smokers are those who smoke on average less than 10 cigarettes per day. Moderate smokers smoke at least 10 but fewer than 20 cigarettes per day. Heavy smokers are those who smoke 20 or more cigarettes per day.


Download chart**XLS** [XLS format](#)

(27.5 Kb)

Within the group of cigarette smokers the rate of self-reported LSI varied. Higher consumption levels were associated with increased likelihood of having reported an LSI. Nearly a third (31%) of light smokers (up to 10 cigarettes per day) reported having an LSI, about the same as the proportion of those who had never smoked cigarettes (32%). This increased to 44% among heavy smokers (20 or more cigarettes per day).

Regional self-reported long-standing illness or disability

Rates of self-reported LSI and limiting LSI were higher in Wales than in England or Scotland. More than one in four adults in Wales (27%) reported having a limiting LSI, compared with around one in five people in England (19%) and Scotland (20%). This remained the case once the effects of factors such as age and employment status had been removed. There was some variation in the rates across England, with rates tending to be higher in the North East (23%). Rates of self-reported limiting LSIs were lowest in the South East (17%), London (18%), and East of England (17%), Figure 7.

Figure 7: Regional self-reported long-standing illness or disability, Great Britain, 2013

Source: Opinions and Lifestyle Survey - Office for National Statistics

Download chart

[XLS](#) [XLS format](#)
(28.5 Kb)


Effects of health conditions lasting 12 months or more

The most commonly reported difficulties caused by long-term health conditions lasting for 12 months or more were issues with mobility (13%), stamina, breathing and fatigue (10%) and dexterity (7%), Figure 8.

Self-reported memory difficulties and physical health conditions increased with age. This was not the case for difficulties with social skills, behaviour, learning and ability to understand or

concentrate. People between the ages of 25 and 64 reported mental health difficulties more frequently (5%) than those in younger (3%) or older (2%) age groups. This reflects findings in the [Welsh Health Survey 2013](#) and the [Scottish Health Survey 2013](#) that physical illness increases with age and mental wellbeing tends to be highest in the youngest and oldest age groups.

There were differences between men and women in the reported effects of long-term health conditions. Men reported more hearing difficulties than women (5% vs. 4%), while women reported more difficulties with dexterity (8% vs. 6%) and mobility (14% vs. 11%) than men.

Figure 8: Effects of health conditions lasting 12 months or longer, Great Britain, 2013

Source: Opinions and Lifestyle Survey - Office for National Statistics

Notes:

1. This include health conditions that are expected to last for 12 months or longer if they have not already done so.

Download chart

XLS [XLS format](#)

(28 Kb)

Background notes**1. The Opinions and Lifestyle Survey**

The data in this report were collected on the [Opinions and Lifestyle Survey \(OPN\)](#) - an omnibus survey run by the Office for National Statistics. Each survey wave lasts for one month and is open for both government and non-government organisations to run questions.

The OPN is currently the only randomised probability sample omnibus survey in Great Britain and provides a fast, reliable and flexible service to customers.

More information on the survey and survey methodology can be found in the [Opinions and Lifestyle Survey Information Guide \(175.5 Kb Pdf\)](#).

2. How to commission a module on the survey

Clients can enquire about purchasing modules of questions by emailing the survey manager at opinions@ons.gsi.gov.uk.

3. Comparability

This report provides information on the health of adults in Great Britain, and follows on from the series of releases from the General Household Survey (GHS) and General Lifestyle Survey (GLF).

The OPN and GHS/GLF provide comparable results. However there are some differences in the design and content of the surveys. More information can be found in the '[Opinions and Lifestyle Survey, Smoking Habits Amongst Adults, 2012](#)' publication.

4. Coherence

There are a number of other sources of long-standing illness and disability data. These have been listed below, together with a brief explanation of the comparability with the OPN.

Health Survey for England (Health and Social Care Information Centre), Welsh Health Survey (Welsh Government) and Scottish Health Survey (Scottish Government)

There are some differences in the questions asked about health between these surveys. In the Welsh Health Survey respondents are asked whether they are currently being treated for any of a range of illnesses. They are also asked whether their daily activities are limited by a health problem or disability lasting (or expected to last) at least 12 months.

In the Scottish Health Survey and the Health Survey for England respondents are asked if they have any physical or mental health condition or illness lasting, or likely to last, for twelve months or more. Those who report having a condition are then asked whether it limits their daily activities. The questions are based on self-perception of illness rather than doctor diagnosis.

5. Reliability

Self-reports of long-standing illness or disability may differ from rates of diagnosed long-standing illness or disability due to differences in the way people perceive their health.

6. Approach to statistical significance

Where estimates for different populations have been described as different throughout this commentary, they have been tested and found to be significantly different at 5% significance level ($p < 0.05$).

95% confidence intervals for each table value have been supplied as a [separate table \(78.5 Kb Excel sheet\)](#). Where historical data have been provided, confidence intervals have been supplied for the last two years (2012 and 2013).

7. Details of the policy governing the release of new data are available by visiting www.statisticsauthority.gov.uk/assessment/code-of-practice/index.html or from the Media Relations Office email: media.relations@ons.gsi.gov.uk

The United Kingdom Statistics Authority has designated these statistics as National Statistics, in accordance with the Statistics and Registration Service Act 2007 and signifying compliance with the Code of Practice for Official Statistics.

Designation can be broadly interpreted to mean that the statistics:

- meet identified user needs;
- are well explained and readily accessible;
- are produced according to sound methods; and
- are managed impartially and objectively in the public interest.

Once statistics have been designated as National Statistics it is a statutory requirement that the Code of Practice shall continue to be observed.

Copyright

© Crown copyright 2015

You may use or re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit

www.nationalarchives.gov.uk/doc/open-government-licence/ or write to the Information Policy Team, The National Archives, Kew, London TW9 4DU, or email: psi@nationalarchives.gsi.gov.uk.

This document is also available on our website at www.ons.gov.uk.

Statistical contacts

Name	Phone	Department	Email
Dr Craig Orchard	+44 (0)1633 455755	Office for National Statistics	socialsurveys@ons.gsi.gov.uk

Issuing Body:

Office for National Statistics

Media Contact Details:

Telephone: 0845 604 1858
(8.30am-5.30pm Weekdays)

Emergency out of hours (limited service): 07867 906553

Email:

media.relations@ons.gsi.gov.uk

Table	Title
1	<u>Self-Perception of General Health, 2005-2013</u>
2	<u>Self-Reported Prevalence of Long-Standing Illness or Disability, 2005-2013</u>
3	<u>Effects of Physical or Mental Health Conditions Lasting or Expected to Last for 12 Months or More, 2013</u>
4	<u>Prevalence of Long-Standing Illness or Disability, by Socio-Economic Classification, 2013</u>
5	<u>Prevalence of Long-Standing Illness or Disability, by Employment Status, 2013</u>
6	<u>Gross Personal Income and Prevalence of Long-Standing Illness or Disability, 2013</u>
7	<u>Relationship Status and Prevalence of Long-Standing Illness or Disability, 2013</u>
8	<u>Prevalence of Long-Standing Illness or Disability, by Region, 2013</u>
9	<u>Prevalence of Long-Standing Illness or Disability, by Smoking Status, 2013</u>

1 Self-Perception of General Health, 2005-2013

All persons aged 16 and over, Great Britain					Percentages
	Very Good	Good	Fair	Bad	Very Bad
2005 ^{1 2 3}	34	41	18	6	1
2006 ^{2 3 4}	33	43	17	5	1
2007 ⁴	34	43	17	5	1
2008 ⁴	39	40	16	4	1
2009 ⁴	40	39	15	5	1
2010 ⁴	39	40	15	5	1
2011 ⁴	34	43	18	5	1
2012	39	40	15	4	1
2013	38	41	15	4	1
Weighted base 2013 (000s) = 100%	48,822	48,822	48,822	48,822	48,822
Unweighted sample 2013 ⁵	12,010	12,010	12,010	12,010	12,010

Source: Opinions and Lifestyle Survey, General Lifestyle Survey and General Household Survey;
Office for National Statistics

- 1 Estimates prior to 2006 cover a fiscal year rather than a calendar year.
- 2 Estimates for 2005 and 2006 both include the period 1 January 2006 to 31 March 2006 as a result of change in survey year from a fiscal year to a calendar year.
- 3 There are potential exposure and order effects associated with the five category general health question prior to 2008. Particular care should therefore be taken when trying to make conclusions between periods before 2008 and those from 2008 onwards.
- 4 Estimates for 2006 to 2011 include longitudinal data.
- 5 All unweighted samples are rounded to the nearest 10.

2 Self-Reported Prevalence of Long-Standing Illness or Disability, By Age and Gender 2005-2013

All persons aged 16 and over

Percentages

	Men						Women						All persons					
	16 to 24	25 to 44	45 to 64	65 to 74	75 and over	All aged 16 and over	16 to 24	25 to 44	45 to 64	65 to 74	75 and over	All aged 16 and over	16 to 24	25 to 44	45 to 64	65 to 74	75 and over	All aged 16 and over
Proportion with long-standing illness or disability																		
2005 ^{1 2}	15	24	44	57	64	36	17	26	43	61	64	38	16	25	44	59	64	37
2006 ^{2 3}	15	23	45	63	70	37	16	26	44	64	70	39	16	24	45	63	70	38
2007 ³	13	22	43	60	66	36	17	23	41	56	65	36	15	23	42	58	65	36
2008 ³	12	20	39	58	64	34	16	21	38	53	68	35	14	20	39	56	67	34
2009 ³	12	20	42	60	64	35	16	22	39	56	67	36	14	21	40	58	66	35
2010 ³	12	19	41	58	68	34	17	23	40	55	68	36	15	21	41	56	68	35
2011 ³	13	22	42	59	69	36	21	26	41	57	67	39	17	24	42	58	68	38
2012	14	20	38	56	69	33	15	22	39	57	66	35	14	21	39	57	67	34
2013	14	19	42	59	68	34	17	22	42	55	70	37	15	21	42	57	69	36
Weighted base 2013 (000s)	3,415	8,157	7,631	2,746	1,964	23,912	3,298	8,093	7,876	3,001	2,623	24,890	6,713	16,250	15,507	5,746	4,586	48,802
Unweighted sample 2013	420	1,550	1,850	910	650	5,370	490	2,040	2,110	1,100	910	6,640	910	3,590	3,960	2,000	1,560	12,010
Proportion with limiting long-standing illness or disability																		
2005 ^{1 2}	7	13	26	35	44	21	9	14	26	38	48	23	8	14	26	37	46	22
2006 ^{2 3}	6	11	23	37	47	20	8	13	27	39	51	24	7	12	25	38	50	22
2007 ³	6	12	25	37	47	21	9	13	25	36	48	23	7	13	25	37	48	22
2008 ³	6	10	22	33	45	19	10	11	23	34	48	21	8	11	23	33	46	20
2009 ³	7	10	25	36	42	21	8	13	23	35	50	22	7	12	24	36	47	21
2010 ³	7	11	23	34	45	20	12	13	23	34	50	23	9	12	23	34	48	21
2011 ³	5	12	24	35	44	21	13	14	25	37	48	24	9	13	25	36	47	23
2012	7	10	20	31	48	18	6	11	23	31	47	20	6	10	21	31	48	19
2013	8	9	21	33	45	19	7	11	23	31	49	21	8	10	22	32	47	20
Weighted base 2013 (000s)	3,415	8,157	7,631	2,746	1,964	23,912	3,298	8,093	7,876	3,001	2,623	24,890	6,713	16,250	15,507	5,746	4,586	48,802
Unweighted sample 2013 ⁴	420	1,550	1,850	910	650	5,370	490	2,040	2,110	1,100	910	6,640	910	3,590	3,960	2,000	1,560	12,010

Source: Opinions and Lifestyle Survey, General Lifestyle Survey and General Household Survey; Office for National Statistics

1 Estimates prior to 2006 cover a fiscal year rather than a calendar year.

2 Estimates for 2005 and 2006 both include the period 1 January 2006 to 31 March 2006 as a result of a change in survey year from a fiscal year to a calendar year.

3 Estimates for 2006 to 2011 include longitudinal data.

4 All unweighted samples are rounded to the nearest 10.

3 Effects of Physical and Mental Health Conditions Lasting for 12 Months or More, 2013^{1 2}

All persons aged 16 and over

Percentages

	Men						Women						All persons					
	16 to 24	25 to 44	45 to 64	65 to 74	75 and over	All aged 16 and over	16 to 24	25 to 44	45 to 64	65 to 74	75 and over	All aged 16 and over	16 to 24	25 to 44	45 to 64	65 to 74	75 and over	All aged 16 and over
Vision	1	2	4	7	15	4	1	2	4	6	17	4	1	2	4	7	16	4
Hearing	1	1	4	12	25	5	0	1	3	6	18	4	0	1	4	9	21	5
Mobility	3	4	12	22	36	11	1	5	15	23	47	14	2	5	13	22	42	13
Dexterity	1	3	7	10	18	6	1	3	9	12	23	8	1	3	8	11	21	7
Learning or ability to understand or concentrate	4	2	3	2	6	3	3	2	2	2	2	2	3	2	3	2	4	3
Memory	1	2	4	6	15	4	1	2	4	5	10	4	1	2	4	5	12	4
Mental Health	2	4	5	2	2	4	4	6	6	3	3	5	3	5	6	2	2	4
Stamina, breathing or susceptibility to fatigue	4	4	10	18	25	9	5	5	12	14	25	10	4	4	11	16	25	10
Social skills or behaviour	2	1	1	0	0	1	2	0	0	0	0	1	2	1	1	0	0	1
Any other issues	3	4	6	8	8	5	2	5	5	9	6	5	3	5	6	9	7	5
Weighted base (000s)	3,431	8,261	7,662	2,748	1,981	24,083	3,317	8,190	7,969	3,001	2,655	25,130	6,748	16,451	15,630	5,749	4,636	49,214
Unweighted sample 2013 ³	420	1,560	1,850	910	660	5,390	490	2,060	2,130	1,100	910	6,690	910	3,630	3,980	2,000	1,570	12,080

Source: Opinions and Lifestyle Survey, General Lifestyle Survey and General Household Survey; Office for National Statistics

1 Includes physical and mental health conditions that are expected to last for 12 months or more.

2 People may have health conditions that affect them in more than one of the given areas.

3 All unweighted samples are rounded to the nearest 10.

4 Prevalence of Long-Standing Illness or Disability, by Socio-Economic Classification, 2013

All persons aged 16 and over			Percentages
	Managerial and professional occupations	Intermediate occupations	Routine and manual occupations
Proportion with a long-standing illness or disability	27	29	32
Proportion with a limiting long-standing illness or disability	11	12	16
Weighted bases (000s) = 100%			
Long-standing illness or disability	14,242	8,433	11,757
Limiting long-standing illness or disability	14,229	8,433	11,757
Unweighted samples ¹			
Long-standing illness or disability	3,330	1,990	2,770
Limiting long-standing illness or disability	3,320	1,990	2,770

Source: Opinions and Lifestyle Survey, Office for National Statistics

¹ All unweighted samples are rounded to the nearest 10.

5 Prevalence of Long-Standing Illness or Disability, by Employment Status, Great Britain, 2013¹

All persons aged 16 and over

Percentages

	In Employment	ILO Unemployed ²	Economically Inactive ³
Proportion with a long-standing illness or disability	24	33	56
Proportion with a limiting long-standing illness or disability	9	20	37
Weighted bases (000s) = 100%			
Long-standing illness or disability	28,859	2,805	17,138
Limiting long-standing illness or disability	28,847	2,805	17,138
Unweighted samples⁴			
Long-standing illness or disability	6,320	620	5,080
Limiting long-standing illness or disability	6,320	620	5,080

Source: Opinions and Lifestyle Survey, Office for National Statistics

1 International Labour Organisation (ILO) definitions of employment status have been used.

2 Unemployed people are those not currently working, but who are looking for work.

3 Economically inactive people are those not working nor looking for work. This includes retired people and students.

4 All unweighted samples are rounded to the nearest 10.

6 Gross Personal Income and Prevalence of Long-Standing Illness or Disability, 2013

All persons aged 16 and over

Percentages

	Up to £10,000	£10,000 up to £15,000	£15,000 up to £20,000	£20,000 up to £30,000	£30,000 up to £40,000	£40,000 up to £50,000	£50,000 or more
Proportion with a long-standing illness or disability	45	41	34	30	27	25	24
Proportion with a limiting long-standing illness or disability	30	25	17	12	9	10	8
Weighted bases (000s) = 100%							
Long-standing illness or disability	13,026	7,843	5,859	7,179	3,982	1,925	2,967
Limiting long-standing illness or disability	13,026	7,843	5,859	7,179	3,982	1,925	2,955
Unweighted samples ²							
Long-standing illness or disability	3,340	2,180	1,490	1,770	920	450	640
Limiting long-standing illness or disability	3,340	2,180	1,490	1,770	920	450	640

Source: Opinions and Lifestyle Survey, Office for National Statistics

1 Annual gross personal income covers only the income that an individual receives directly, before deductions for tax etc. Household income, to which an individual may have access, is not included.

2 All unweighted samples are rounded to the nearest 10.

7 Relationship Status and Prevalence of Long-Standing Illness or Disability, 2013

All persons aged 16 and over

	Percentages			
	Single	Married ¹	Co-habiting	Widowed/ Divorced/ Separated ²
Proportion with a long-standing illness or disability	26	38	25	54
Proportion with a limiting long-standing illness or disability	15	19	12	35
Weighted bases (000s) = 100%				
Long-standing illness or disability	11,478	24,703	5,671	6,950
Limiting long-standing illness or disability	11,478	24,691	5,671	6,950
Unweighted samples ³				
Long-standing illness or disability	2,480	5,420	1,140	2,980
Limiting long-standing illness or disability	2,480	5,420	1,140	2,980

Source: Opinions and Lifestyle Survey, Office for National Statistics

- 1 Includes those in a same-sex Civil Partnership.
- 2 Includes those who had been in a same-sex Civil Partnership.
- 3 All unweighted samples are rounded to the nearest 10.

8 Prevalence of Long-Standing Illness or Disability, by Region 2013

All persons aged 16 and over

Percentages

	North East	North West	Yorkshire and The Humber	East Midlands	West Midlands	East of England	London	South East	South West	England	Wales	Scotland
Proportion with a long-standing illness or disability	42	37	36	36	35	34	32	34	39	35	42	36
Proportion with a limiting long-standing illness or disability	23	21	21	20	19	17	18	17	20	19	27	20
Weighted bases (000s) = 100%												
Long-standing illness or disability	2,135	5,547	4,268	3,629	4,328	4,756	6,215	6,886	4,323	42,088	2,457	4,256
Limiting long-standing illness or disability	2,135	5,547	4,268	3,629	4,328	4,756	6,215	6,886	4,311	42,076	2,457	4,256
Unweighted samples ¹												
Long-standing illness or disability	550	1,570	1,210	950	1,050	1,160	1,040	1,690	1,070	10,290	630	1,090
Limiting long-standing illness or disability	550	1,570	1,210	950	1,050	1,160	1,040	1,690	1,070	10,290	630	1,090

Source: Opinions and Lifestyle Survey, Office for National Statistics

¹ All unweighted samples are rounded to the nearest 10.

9 Prevalence of Long-Standing Illness or Disability, by Smoking Status 2013

All persons aged 16 and over

	Percentages						
	Never Smoked	Ex- Smokers	All Non- Smokers	Light Smokers ¹	Moderate Smokers ²	Heavy Smokers ³	All Smokers
Proportion with a long-standing illness or disability	32	45	35	31	38	44	37
Proportion with a limiting long-standing illness or disability	16	25	19	19	25	31	24
Weighted bases (000s) = 100%							
Long-standing illness or disability	28,266	11,172	39,445	3,391	3,906	1,999	9,343
Limiting long-standing illness or disability	28,253	11,172	39,433	3,391	3,906	1,999	9,343
Unweighted samples ⁴							
Long-standing illness or disability	6,580	3,060	9,640	830	990	540	2,370
Limiting long-standing illness or disability	6,580	3,060	9,640	830	990	540	2,370

Source: Opinions and Lifestyle Survey, Office for National Statistics

1 Up to 10 cigarettes per day.

2 10 up to 20 cigarettes per day.

3 20 or more cigarettes per day.

4 All unweighted samples are rounded to the nearest 10.